

A new diagnosis of asthma or COPD is linked to smoking cessation – the Tromsø study

Signe Elise Danielsen¹
Maja-Lisa Løchen¹
Astri Medbø¹
Monica Linea Vold²
Hasse Melbye³

¹Department of Community Medicine, The Arctic University of Norway,

²Department of Respiratory Medicine, University Hospital of North Norway,

³General Practice Research Unit, Department of Community Medicine, UiT The Arctic University of Norway, Tromsø, Norway

Background: Patients with COPD have had a lower tendency to quit smoking compared to patients with coronary heart disease (CHD). We wanted to investigate if this is still true in a Norwegian population.

Methods: Our data came from the fifth and sixth Tromsø surveys, which took place in 2001–2002 and 2007–2008. The predictors of smoking cessation were evaluated in a cohort of 4,497 participants who had stated their smoking status in both surveys.

Results: Of the 4,497 subjects in the cohort, 1,150 (25.6%) reported daily smoking in Tromsø 5. In Tromsø 6, 428 had quit (37.2%). A new diagnosis of obstructive lung disease (asthma or COPD) and CHD were both associated with increased quitting rates, 50.6% ($P=0.01$) and 52.1% ($P=0.02$), respectively. In multivariable logistic regression analysis with smoking cessation as outcome, the odds ratios (ORs) of a new diagnosis of obstructive lung disease and of CHD were 1.7 (1.1–2.7) and 1.7 (1.0–2.9), respectively. Male sex had an OR of 1.4 (1.1–1.8) compared to women in the multivariable model, whereas the ORs of an educational length of 13–16 years and ≥ 17 years compared to shorter education were 1.6 (1.1–2.2) and 2.5 (1.5–4.1), respectively.

Conclusion: The general trend of smoking cessation in the population was confirmed. Increased rates of smoking cessation were associated with a new diagnosis of heart or lung disease, and obstructive lung disease was just as strongly linked to smoking cessation as was CHD. This should encourage the pursuit of early diagnosis of COPD.

Keywords: smoking cessation, cohort study, COPD, asthma, coronary heart disease

Background

Smoking cessation is crucial for a better prognosis in patients with coronary heart disease (CHD)^{1–3} and COPD.^{2,4,5} Acute heart attacks are associated with high smoking cessation rates. In patients with a first myocardial infarction (MI) in the 1970s,³ among whom 78% were smoking, the quitting rate was 55%. In a systematic review from 2003, the mean rate of smoking cessation within a year after a new diagnosis of CHD was found to be 45%.¹ A lower quitting rate was observed in patients with COPD, in particular in subjects not participating in smoking cessation programs. Among smoking COPD patients in a Swedish study, 10% of those receiving usual care had stopped smoking after 3 years, compared to 38% of those who took part in smoking cessation groups.⁶ Among hospitalized patients with respiratory and cardiac diseases who took part in an intensive smoking cessation program in Singapore, 60% of the cardiac patients and 40% of the respiratory patients were still abstinent from smoking after 2 months.⁷

The mortality due to CHD has decreased considerably in the Western world in the last few decades.⁸ Evidence shows that decreased smoking prevalence in the general population, particularly among patients with heart disease, is an important reason for

Correspondence: Hasse Melbye
General Practice Research Unit,
MH-building, Department of Community
Medicine, UiT The Arctic University of
Norway, 9037 Tromsø, Norway
Email hasse.melbye@uit.no

this decrease.^{8–11} The mortality due to COPD has decreased less.¹² A stronger tendency of persistent smoking among COPD patients may partly explain this difference.

COPD patients' reluctance to quit smoking may lead to a feeling of hopelessness among the doctors who treat them.¹³ Our aim was to determine how the quitting rate in adults is influenced by getting a diagnosis of obstructive lung disease in the population-based Tromsø study. We also wanted to investigate whether influence from this life event differs from being diagnosed with CHD.

Methods

Subjects

A large proportion of the adult population in Tromsø has participated in health surveys (the Tromsø study) since 1974. Tromsø is a university city in northern Norway with ~72,600 inhabitants. The Tromsø study so far consists of six surveys, and this study deals with subjects who participated in both Tromsø 5 (2001–2002) and Tromsø 6 (2007–2008). The attendance rates (of those invited) in Tromsø 5 and Tromsø 6 were 79% and 66%, respectively. Details of the participants have previously been described.¹⁴ We included participants who answered the questions on smoking in both surveys. No smoking cessation support was offered during the surveys.

The Regional committee for Medical and Health Research Ethics and the Norwegian Data Inspectorate approved the Tromsø 5 and Tromsø 6 surveys with a license for further analysis on non-identifiable data, like this study. All the participants gave written informed consent.

Examinations

In both surveys, the invitation included a questionnaire with questions on smoking habits, education, and diseases. We classified subjects reporting daily smoking in Tromsø 5, but not in Tromsø 6, as quitters. We divided length of education into three categories, ≤12 years, 13–16 years, and ≥17 years. Table 1 shows the reported diseases. For each disease in the questionnaire the participants were asked whether they have or have had the disease in question. We classified diseases reported in Tromsø 6 and not in Tromsø 5 as new diagnoses. We categorized MI and angina pectoris as “CHD” and asthma and COPD as “obstructive lung disease”.

Statistical analysis

The frequencies of current smoking at baseline (in Tromsø 5) and smoking cessation among smokers were calculated by subject characteristics, and the statistical differences between subgroups were analyzed by chi-square statistics. We analyzed

Table 1 Daily smoking according to characteristics of the cohort of 4,497 participants in Tromsø 5 (2001–2002) who also took part in Tromsø 6 (2007–2008)

	n	Smoking, n (%)	P-value ^a
Sex			
Men	1,852	462 (24.9)	0.4
Women	2,645	688 (26.0)	
Age (years)			
30–49	671	233 (34.7)	<0.001 ^b
50–69	2,879	733 (26.8)	
70–81	947	144 (15.2)	
Education			
0–12 years	3,265	869 (26.6)	<0.001 ^b
13–16 years	691	168 (24.3)	
>16 years	407	75 (18.4)	
Not answered	134	38 (28.3)	
Self-reported diseases			
Asthma			
Yes	366	91 (24.9)	0.7
No	4,048	1,040 (25.7)	
Not answered	85	19 (22.3)	
COPD			
Yes	192	84 (43.8)	<0.001
No	4,201	1,036 (24.1)	
Not answered	104	30 (28.8)	
Asthma or COPD			
Yes	482	145 (30.1)	0.02
No	3,923	982 (25.1)	
Not answered	94	23 (24.5)	
Angina pectoris			
Yes	314	55 (17.5)	0.001
No	4,078	1,072 (26.3)	
Not answered	105	23 (25.3)	
MI			
Yes	238	53 (22.3)	0.2
No	4,168	1,076 (25.8)	
Not answered	91	21 (23.1)	
CHD (angina or infarction)			
Yes	442	84 (19.0)	0.001
No	3,962	1,046 (26.4)	
Not answered	93	20 (21.5)	
Diabetes			
Yes	137	30 (21.9)	0.3
No	4,275	1,099 (25.7)	
Not answered	85	21 (24.7)	
Stroke			
Yes	105	19 (18.1)	0.08
No	4,280	1,103 (25.8)	
Not answered	112	28 (25.0)	

Notes: ^aStatistical significance of difference between subgroups, “not answered” excluded. ^bChi-square trend.

Abbreviations: MI, myocardial infarction; CHD, coronary heart disease.

predictors of smoking cessation by binary logistic regression. Significant variables in univariable analysis ($P < 0.1$), as well as age and sex, were entered in the multivariable analysis. For the statistical analyses, we used the IBM SPSS statistic, Version 21 (IBM Corporation, Armonk, NY, USA).

Results

A total of 4,631 subjects participated in both the fifth and the sixth surveys of the Tromsø study, and 4,497 subjects comprising 1,852 men and 2,645 women answered the questions on smoking in both surveys. The mean time difference between attending the two surveys was 6.2 years (standard deviation 0.5 years). The mean age at Tromsø 5 was 61.1 years (range 30–81 years); 1,150 (25.6%) reported daily smoking and 1,753 (39.0%) reported former smoking. The highest prevalence of smoking (43.8%) was found in subjects with self-reported COPD, whereas the frequency among those with self-reported CHD was 19.0%. The characteristics of the population in Tromsø 5 and the respective frequencies of smoking are listed in Table 1.

In the sixth survey, 765 subjects (17.0%) reported current smoking, and 428 of the 1,150 who had been smoking in Tromsø 5 had quit (37.2%). A significantly higher quitting rate was found in men compared to women, and the quitting rate increased with the length of education. Self-reported diseases registered in Tromsø 5 had no significant impact on the quitting rate (Table 2). The number of days between attending the two surveys had no impact on the quitting rate (odds ratio =1, $P=0.8$).

Table 3 shows the frequencies of new self-reported diseases reported in Tromsø 6, but not in Tromsø 5, and the association between a new diagnosis and smoking cessation. New diagnoses of asthma/COPD ($n=79$) and CHD ($n=73$) were both associated with increased frequency of quitting, 50.6% ($P=0.01$) and 52.1% ($P=0.008$), respectively. A new diagnosis of asthma/COPD or CHD was reported in 144 subjects. Both men and women belonging to any of these diagnostic groups showed increased quitting rates, 54.9% vs 39.8% ($P=0.02$) in men and 47.1% vs 32.7% ($P=0.009$) in women. The impact of getting these new diagnoses was particularly strong among the subjects with education length no longer than 12 years ($P=0.001$; Figure 1). Subjects with education length up to 12 years were also more frequently diagnosed with these diseases compared to those with higher education, 16.0% and 9.5% ($P=0.01$), respectively. Among subjects with higher education, a high quitting rate was independent of a new diagnosis (Figure 1).

A new diagnosis of both CHD and obstructive lung disease was significantly associated with smoking cessation when analyzed by logistic regression and also after adjusting for sex and education level in multivariable logistic regression analyses (Table 4).

Table 2 Smoking cessation between 2001–2002 (Tromsø 5) and 2007–2008 (Tromsø 6) according to characteristics of 1,150 smoking participants in Tromsø 5

	n	Quit smoking, n (%)	P-value ^a
Sex			
Men	462	194 (42.0)	0.006
Women	688	234 (34.0)	
Age (years)			
30–49	233	87 (37.7)	0.6 ^b
50–69	773	292 (37.8)	
70–81	144	49 (34.0)	
Education			
0–12 years	869	300 (34.5)	<0.001 ^b
13–16 years	168	74 (44.0)	
>16 years	75	42 (56.0)	
Not answered	38	12 (31.6)	
Self-reported diseases			
Asthma			
Yes	91	26 (28.6)	0.8
No	1,040	394 (37.9)	
Not answered	19	6 (31.6)	
COPD			
Yes	84	35 (41.7)	0.4
No	1,036	384 (37.1)	
Not answered	30	9 (30.0)	
Asthma or COPD			
Yes	145	51 (35.2)	0.5
No	982	371 (37.8)	
Not answered	23	6 (26.1)	
Angina pectoris			
Yes	55	17 (30.9)	0.3
No	1,072	402 (37.5)	
Not answered	23	9 (39.1)	
MI			
Yes	53	17 (32.1)	0.4
No	1,076	403 (37.5)	
Not answered	21	8 (38.1)	
CHD (angina or infarction)			
Yes	84	28 (33.3)	0.3
No	1,046	392 (37.5)	
Not answered	20	8 (40.0)	
Diabetes			
Yes	30	7 (23.3)	0.1
No	1,099	412 (37.5)	
Not answered	21	9 (42.9)	
Stroke			
Yes	19	6 (31.6)	0.6
No	1,103	412 (37.4)	
Not answered	28	10 (35.7)	

Notes: ^aStatistical significance of difference between subgroups, “not answered” excluded. ^bChi-square trend.

Abbreviations: MI, myocardial infarction; CHD, coronary heart disease.

Discussion

Main findings

Half of the participants with a new diagnosis of asthma/COPD or angina/MI quit smoking during the 6 years of follow-up. However, the association between these new diagnoses and smoking cessation was found only in participants

Table 3 Smoking cessation between 2001–2002 (Tromsø 5) and 2007–2008 (Tromsø 6) according to new diagnosis in the same period

	n	Quit smoking, n (%)	P-value ^a
New diagnosis			
Asthma			
Yes	42	20 (47.6)	0.2
No	967	365 (37.7)	
COPD			
Yes	75	36 (48.0)	0.05
No	933	342 (36.7)	
Asthma or COPD			
Yes	79	40 (50.6)	0.01
No	903	331 (36.7)	
Angina pectoris			
Yes	33	15 (45.5)	0.3
No	1,017	379 (37.3)	
MI			
Yes	58	30 (51.7)	0.02
No	995	365 (36.7)	
CHD (angina or infarction)			
Yes	73	38 (52.1)	0.008
No	955	348 (36.4)	
Diabetes			
Yes	33	12 (34.4)	0.9
No	1,035	391 (37.8)	
Stroke			
Yes	37	12 (32.4)	0.5
No	1,037	389 (37.5)	

Notes: ^aStatistical significance of difference between participants with a new (possibly combined) diagnosis and participants without a new (possibly combined) diagnosis in Tromsø 5. In each analysis, participants who reported the diagnosis in question in Tromsø 5, and those who have not answered the question on the diagnosis in any of the surveys, have been excluded.

Abbreviations: MI, myocardial infarction; CHD, coronary heart disease.

with education length up to 12 years. The much higher smoking rate among those with asthma and, in particular, those with COPD at baseline than in those with CHD indicates a shift in attitude toward smoking cessation among subjects with COPD.

Strengths and limitations

The high number of participants and the high attendance rates among those invited are strengths of the study. The prevalences of daily smoking among all the attendees in Tromsø 5 and Tromsø 6 were 31% and 22%, respectively,¹¹ which are close to the national prevalences of 29% and 21%, as registered by Statistics Norway (<https://www.ssb.no>)¹⁵ in 2002 and 2008, respectively. In our study sample, the prevalence was lower at both points of time, 25.6% and 17.0%, respectively, and a healthy survivor effect may have contributed to the low frequency of daily smoking in our subsample. The study is based on questionnaires and not on objective measurements of smoking, such as cotinine and thiocyanate. In a previous Norwegian study, self-reported




Figure 1 Percentage of participants who quit smoking between 2001–2002 (Tromsø 5) and 2007–2008 (Tromsø 6) among 1,064 participants, by education length, and heart and lung disease.

smoking was strongly related to serum thiocyanate if the question was asked in a neutral setting.¹⁶ Although the questions on smoking were included in a self-administered questionnaire, underreporting of daily smoking among those with a new diagnosis cannot be ruled out.

All diagnoses were self-reported, and their correctness could not be confirmed. Shift in diagnosis based on the same illness from Tromsø 5 to Tromsø 6, for instance, between asthma and COPD, may have taken place in some subjects. Such a change has been counted as a new diagnosis. This is no longer a problem when asthma and COPD are combined into one category (asthma and/or COPD) and subjects with a new diagnosis of any of these diseases are compared with those with neither of the diagnoses at both Tromsø 5 and Tromsø 6.

Although we have found associations between a new diagnosis and smoking cessation, we do not know for sure whether the extra cessations in the subgroups with a new diagnosis were really preceded by a new diagnosis. Subjects in these subgroups could have stopped smoking before the diagnosis was given. In these cases, it is likely that symptoms from a pulmonary or heart disease had raised their awareness of the risk of continued smoking.

Comparisons with previous studies

No previous study has, to our knowledge, described smoking cessation in relation to a new diagnosis of heart or lung disease. Most previous studies evaluated smoking cessation

Table 4 ORs for smoking cessation between Tromsø 5 (2001–2002) and Tromsø 6 (2007–2008) according to characteristics of 1,024 smoking participants in Tromsø 5 and new diagnosis reported in Tromsø 6

	Univariable		Multivariable	
	OR (95% CI)	P-value	OR (95% CI)	P-value
Age	1.0 (1.0–1.0)	0.9	1.0 (1.0–1.0)	0.6
Male sex	1.4 (1.1–1.8)	0.01	1.4 (1.1–1.8)	0.02
Education (years)				
13–16	1.5 (1.0–2.1)	0.03	1.6 (1.1–2.2)	0.02
≥17	2.3 (1.4–3.8)	0.001	2.5 (1.5–4.1)	<0.001
New diagnosis of CHD	1.7 (1.1–2.8)	0.02	1.7 (1.0–2.9)	0.04
New diagnosis of asthma or COPD	1.7 (1.1–2.6)	0.02	1.7 (1.1–2.7)	0.02

Abbreviations: OR, odds ratio; CHD, coronary heart disease.

programs, and control groups in such studies can be compared with our participants. However, the quitting rate of 41.7% among subjects with a COPD diagnosis reported in Tromsø 5 was more similar to the rate of 38% among the COPD patients, who had taken part in the smoking cessation program in a Swedish study, than the rate of 10% among the COPD patients on usual care.⁶ In a study from primary care, where smokers received smoking cessation advice and were followed up annually with spirometry, higher rates of abstinence were found among those with COPD than among those with normal lung function.¹⁷ This result is in line with our findings. We found that the quitting rate increased with increasing length of education, and this association is well known from previous studies.¹⁸

Qualitative studies have shown that patients with a COPD diagnosis may have several reasons for not quitting despite the knowledge of harming themselves,¹⁹ and they do not always believe that quitting would give them a better life.²⁰ COPD patients often show little interest in receiving help from medication and describe unassisted quitting as the best method to stop smoking, based on willpower, strong motivation, and internal strength.²¹ In our study, a new diagnosis of COPD or CHD might have given many participants the motivation they needed to quit without assistance. In another qualitative study, the interviewees who had stopped smoking emphasized that persons close to them had a strong influence on their decision to quit.²² This gives a reason to believe that the decreasing acceptance of smoking in the society also influences COPD patients.

Conclusion

It has been suggested that a hard core of smokers will constitute an increasing proportion of COPD patients who still smoke.²³ The high quitting rates among subjects with both an established and a new diagnosis of COPD in our study contradicts this pessimism. The study supports pursuit of early

diagnosis of COPD and gives reasons for a more optimistic attitude among health workers when they discuss smoking cessation with their COPD patients.

Acknowledgment

The authors would like to thank the participants of the Tromsø study.

Author contributions

All authors contributed toward data analysis, drafting and revising the paper and agree to be accountable for all aspects of the work.

Disclosure

The authors report no conflicts of interest in this work.

References

- Critchley JA, Capewell S. Mortality risk reduction associated with smoking cessation in patients with coronary heart disease: a systematic review. *JAMA*. 2003;290(1):86–97.
- Thun MJ, Carter BD, Feskanich D, et al. 50-year trends in smoking-related mortality in the United States. *N Engl J Med*. 2013;368(4):351–364.
- Aberg A, Bergstrand R, Johansson S, et al. Cessation of smoking after myocardial infarction. Effects on mortality after 10 years. *Br Heart J*. 1983; 49(5):416–422.
- Anthonisen NR, Connett JE, Murray RP. Smoking and lung function of Lung Health Study participants after 11 years. *Am J Respir Crit Care Med*. 2002;166(5):675–679.
- Anthonisen NR, Skeans MA, Wise RA, et al. The effects of a smoking cessation intervention on 14.5-year mortality: a randomized clinical trial. *Ann Intern Med*. 2005;142(4):233–239.
- Sundblad BM, Larsson K, Nathell L. High rate of smoking abstinence in COPD patients: smoking cessation by hospitalization. *Nicotine Tob Res*. 2008;10(5):883–890.
- Ong KC, Cheong GN, Prabhakaran L, Earnest A. Predictors of success in smoking cessation among hospitalized patients. *Respirology*. 2005;10(1): 63–69.
- Dalen JE, Alpert JS, Goldberg RJ, Weinstein RS. The epidemic of the 20(th) century: coronary heart disease. *Am J Med*. 2014;127(9):807–812.
- Aspelund T, Gudnason V, Magnusdottir BT, et al. Analysing the large decline in coronary heart disease mortality in the Icelandic population aged 25–74 between the years 1981 and 2006. *PLoS One*. 2010; 5(11):e13957.

10. Young F, Capewell S, Ford ES, Critchley JA. Coronary mortality declines in the U.S. between 1980 and 2000 quantifying the contributions from primary and secondary prevention. *Am J Prev Med.* 2010; 39(3):228–234.
11. Mannsverk J, Wilsgaard T, Mathiesen EB, et al. Trends in modifiable risk factors are associated with declining incidence of hospitalized and nonhospitalized acute coronary heart disease in a population. *Circulation.* 2016;133(1):74–81.
12. Burney PG, Patel J, Newson R, Minelli C, Naghavi M. Global and regional trends in COPD mortality, 1990–2010. *Eur Respir J.* 2015;45(5): 1239–1247.
13. Risør MB, Spigt M, Iversen R, et al. The complexity of managing COPD exacerbations: a grounded theory study of European general practice. *BMJ Open.* 2013;3(12):e003861.
14. Vold ML, Aasebo U, Melbye H. Low FEV1, smoking history, and obesity are factors associated with oxygen saturation decrease in an adult population cohort. *Int J Chron Obstruct Pulmon Dis.* 2014;9: 1225–1233.
15. Statistics Norway. Available from: <https://www.ssb.no/en/helse/statistikker/royk>. Accessed May 31, 2016.
16. Foss OP, Haug K, Hesla PE, Lund-Larsen PG, Vasli LR. Kan vi stole på egenerklærte røykevaner? [Can we rely on self-reported smoking habits?]. *Tidsskr Nor Laegeforen.* 1998;118(14):2165–2168.
17. Stratelis G, Molstad S, Jakobsson P, Zetterstrom O. The impact of repeated spirometry and smoking cessation advice on smokers with mild COPD. *Scand J Prim Health Care.* 2006;24(3):133–139.
18. Goren A, Annunziata K, Schnoll RA, Suaya JA. Smoking cessation and attempted cessation among adults in the United States. *PLoS One.* 2014;9(3):e93014.
19. Eklund BM, Nilsson S, Hedman L, Lindberg I. Why do smokers diagnosed with COPD not quit smoking? – a qualitative study. *Tob Induc Dis.* 2012;10(1):17.
20. van Eerd EA, Risør MB, van Rossem CR, van Schayck OC, Kotz D. Experiences of tobacco smoking and quitting in smokers with and without chronic obstructive pulmonary disease—a qualitative analysis. *BMC Fam Pract.* 2015;16(1):164.
21. Morphet K, Partridge B, Gartner C, et al. Why don't smokers want help to quit? A qualitative study of smokers' attitudes towards assisted vs unassisted quitting. *Int J Environ Res Public Health.* 2015;12(6): 6591–6607.
22. Medbo A, Melbye H, Rudebeck CE. "I did not intend to stop. I just could not stand cigarettes any more." A qualitative interview study of smoking cessation among the elderly. *BMC Fam Pract.* 2011;12:42.
23. Emery S, Gilpin EA, Ake C, Farkas AJ, Pierce JP. Characterizing and identifying "hard-core" smokers: implications for further reducing smoking prevalence. *Am J Public Health.* 2000;90(3):387–394.

International Journal of COPD

Publish your work in this journal

The International Journal of COPD is an international, peer-reviewed journal of therapeutics and pharmacology focusing on concise rapid reporting of clinical studies and reviews in COPD. Special focus is given to the pathophysiological processes underlying the disease, intervention programs, patient focused education, and self management protocols.

Submit your manuscript here: <http://www.dovepress.com/international-journal-of-chronic-obstructive-pulmonary-disease-journal>

Dovepress

This journal is indexed on PubMed Central, MedLine and CAS. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.